

Molecular diagnostics of *Sarcocystis* spp. infections

K. Stojceki, J. Karamon, J. Sroka, T. Cencek

Department of Parasitology and Invasive Diseases, National Veterinary Research Institute,
24-100 Pulawy, Poland

Abstract

Protozoa of the genus *Sarcocystis* (phylum Apicomplexa, family *Sarcocystidae*) is one of the most common parasites affecting animals. Interspecies diagnostic of *Sarcocystis* genus was based on electron microscopy for many years. Because of absence of visible differences between species with reachable magnifications, light microscopy is useless. In many cases serological diagnostic method have lack of sensitivity. A variety of molecular methods have been developed and used to detect and identify *Sarcocystis* spp. and to assess the genetic diversity among this protozoan from different population/hosts. Nowadays, molecular diagnostic is the common, time/cost effective method used all over the world to interspecies differentiation.

Key words: sarcocystis, sarcocystiosis, molecular diagnostics, PCR

Introduction

Protozoa of the genus *Sarcocystis* (phylum Apicomplexa), is one of the most common parasites affecting animals. Until now, about 130 species have been recognized in this genus (Dubey et al. 1989b, Tenter 1995).

The intermediate host (herbivores or omnivores) becomes infected by ingestion of sporulated oocysts or sporocysts. Each sporocyst contains four sporozoites, which are released in the digestive tract. These forms penetrate into blood vessels and multiply as merozoites (Dubey et al. 1989b). Afterwards, merozoites get into the muscles where numerous asexual divisions occur. Finally, mature sarcocyst containing numerous bradyzoites are developed. The specific definitive host becomes infected by ingestion of muscles with mature *Sarcocystis* spp. cyst. Bradyzoites are released in host intestine and penetrate into erythrocytes where sexual reproduction (gametogony) and

oocyst formation occurs. Disintegration of oocysts can occur, and sporocysts may be found in the feces (Fayer 2004).

Sarcocystiosis (sarcocystosis expression is used by many researchers) is a common and cosmopolitan infection among mammals (Table 1), birds, lower vertebrates and humans.

Acute sarcocystiosis in animal intermediate hosts is characterized by encephalitis, inflammation of the brain and spinal cord, bleeding diathesis. It may cause fetal death, premature delivery, abortions in pregnant animals (Tenter 1995, Caspari et al. 2010). Mild and chronic sarcocystiosis leads to a decrease in body weight and amount of fur (Tenter 1995). Moreover, major changes were also observed in the animals behavior (Reiner et al. 2009).

The WHO reported (1981) that, about 50% of parasitic cysts found in muscles of cattle and pigs belong to the species *S. hominis* and *S. suihominis*, respectively (Acha and Szyfres 2003). High prevalence

Table 1. *Sarcocystis* spp. in selected mammals. According to Tenter (1995) with modification of Elsheikha and Mansfield (2007) and Olias et al. (2009).

Intermediate host	Species of parasite	Definitive host
Cattle (<i>Bos taurus</i>)	<i>Sarcocystis cruzi</i>	Canidae, Raccoon (<i>Procyon lotor</i>)
	<i>Sarcocystis hirsuta</i>	Felidae
	<i>Sarcocystis hominis</i>	Primates
Yak (<i>Bos grunniens</i>)	<i>Sarcocystis poephagi</i>	Unknown
	<i>Sarcocystis poephagicanis</i>	Canidae
Goat (<i>Capra aegagrus hircus</i>)	<i>Sarcocystis hircicanis</i>	Dog (<i>Canis familiaris</i>)
	<i>Sarcocystis capracanis</i>	Canidae
	<i>Sarcocystis moulei</i>	Cat (<i>Felis catus</i>)
	<i>Sarcocystis cuprifelis</i>	Cat (<i>Felis catus</i>)
Sheep (<i>Ovis aries</i>)	<i>Sarcocystis arieticanis</i>	Dog (<i>Canis familiaris</i>)
	<i>Sarcocystis tenella</i>	Canidae
	<i>Sarcocystis medusififormis</i>	Cat (<i>Felis catus</i>)
	<i>Sarcocystis gigantea</i>	Cat (<i>Felis catus</i>)
Reindeer (<i>Rangifer tarandus</i>)	<i>Sarcocystis hardangeri</i>	Unknown
	<i>Sarcocystis tarandivulpes</i>	Canidae, Raccoon (<i>Procyon lotor</i>)
	<i>Sarcocystis tarandi</i>	Unknown
	<i>Sarcocystis rangiferi</i>	Unknown
	<i>Sarcocystis rangi</i>	Fox (<i>Vulpes vulpes</i>)
	<i>Sarcocystis grueneri</i>	Canidae, Raccoon (<i>Procyon lotor</i>)
Dromedary (<i>Camelus dromedarius</i>)	<i>Sarcocystis cameli</i>	Dog (<i>Canis familiaris</i>)
Lama (<i>Lama glama</i>)	<i>Sarcocystis aucheniae</i>	Dog (<i>Canis familiaris</i>)
Horse (<i>Equus ferus caballus</i>) and Donkey (<i>Equus africanus asinus</i>)	<i>Sarcocystis bertrami</i>	Dog (<i>Canis familiaris</i>)
	<i>Sarcocystis neurona</i>	Opossum (<i>Monodelphis domestica</i>) (Elsheikha and Mansfield 2007)
	<i>Sarcocystis equicanis</i>	Dog (<i>Canis familiaris</i>)
	<i>Sarcocystis fayeri</i>	Dog (<i>Canis familiaris</i>)
Rabbit (<i>Oryctolagus cuniculus</i>)	<i>Sarcocystis cuniculi</i>	Cat (<i>Felis catus</i>)
Chicken (<i>Gallus gallus domesticus</i>)	<i>Sarcocystis horvathi</i>	Dog (<i>Canis familiaris</i>) (Olias et al. 2009)
Mallard duck (<i>Anas platyrhynchos</i>)	<i>Sarcocystis rileyi</i>	Striped skunks (<i>Mephitis mephitis</i>)
Domestic pig (<i>Sus scrofa</i> f. <i>domestica</i>)	<i>Sarcocystis miescheriana</i>	Canidae, Raccoon (<i>Procyon lotor</i>)
	<i>Sarcocystis suihominis</i>	Primates
	<i>Sarcocystis porcifelis</i>	Cat (<i>Felis catus</i>)
Dog (<i>Canis familiaris</i>)	<i>Sarcocystis canis</i>	Unknown

of muscles sarcocystiosis in pigs was confirmed by data from India, where investigators found that *S. suihominis* and *S. miescheriana* have been determined in 47% and 43% of pigs, respectively (Saleque and Bhatia 1991).

Sarcocystiosis in definitive hosts (Table 1) is most often asymptomatic, however self-limiting mild diarrhea has been observed (Dubey et al. 1989b).

As a result of *Sarcocystis* spp. zoonotic transmission, there are two known infection scenarios. First: one can act as definitive host (gastrointestinal

infection). Second: one can act as intermediate host (muscle infection).

Humans are definitive hosts of two *Sarcocystis* species: *S. hominis* and *S. suihominis* (Dubey et al. 1989b). Some investigators suspect that this type of infection can be connected with abdominal pain, nausea, diarrhea, eosinophilia, inflammatory bowel disease, dyspnoea, increased pulse, and decreased appetite in some patients (Fayer 2004). Lack of hygiene and raw meat consumption are heightening risk factor of gastrointestinal infection for these protozoa

(Wilairatana et al. 1996). Sarcocystiosis connected with gastrointestinal symptoms is present worldwide, as it was confirmed by numerous researches (Dubey et al. 1989b, Fayer 2004). The prevalence of infection was 10% of the adult Laos citizens (Giboda et al. 1991). Among Tibetan citizens, 21.8% were *S. hominis*-positive and 0.06-7% were *S. suihominis*-positive. Also in Malaysia high levels of *Sarcocystis* spp. prevalence (21%) were found (Wong and Pathmanathan 1992). *Sarcocystis* infection was detected in 23.2% of stool samples originated from Thai workers (Wilairatana et al. 1996).

Human muscle sarcocystiosis is seldom detected however, this type of infection has been reported in India, Thailand, Egypt and Malaysia, so far (Acha and Szyfres 2003). In Egypt, 46% among 42 people with rheumatoid disorders examined by western blot, were *Sarcocystis*-positive. Trichinoscopic investigations carried out in 112 Danes showed *Sarcocystis* spp. in muscles of four persons (Greve 1985). The problem of *Sarcocystis* spp. in human muscles remains still unclear. Researchers suspect that this parasite can cause inflammation of muscles, heart diseases, rheumatic pains and abortion (Greve 1985, Pamphlett and O'Donoghue 1990, Habeeb et al. 1996). Moreover, it can be an opportunistic infection in patients with AIDS or immunodeficiency disorders. The species affinity of *Sarcocystis* parasites found in human muscles is not well identified so far. (Arness et al. 1999).

Basic diagnostic methods of *Sarcocystis* spp. are simple and inexpensive. One can use naked eye examination, light microscopy or histology (hematoxylin eosin staining) (Jehle et al. 2009), however, there are some more sophisticated methods, such as: immunofluorescence antibody test (IFAT) (Moré et al. 2011), ELISA (Heckerth and Tenter 1999), Western Blot (Abdul-Rahman et al. 2002). More sensitive serological tests, which could distinguish *S. hominis* from *S. suihominis*, have not yet been developed (Tenter et al. 1991).

Interspecies diagnostic of *Sarcocystis* genus has been based on electron microscopy for many years. Light microscopy is useless, because of absence of visible differences between species with reachable magnifications (Dubey et al. 1989a, b), however, electron microscopy is expensive and time-consuming (Pritt et al. 2008). Thus, molecular biology techniques are very helpful in the diagnosis of this protozoan species.

A variety of molecular methods have been developed and used to detect and identify *Sarcocystis* spp. and to assess the genetic diversity among this protozoan from different population/hosts. Ribosomal DNA sequences are the most common molecular markers used in *Sarcocystis* spp. differentiation (Dahlgren and Gjerde 2008, Rosenthal et al. 2008, Xiang et

al. 2009). This fragment of nucleic acid consists of small ribosomal subunit DNA (SSU rDNA), large ribosomal subunit DNA (LSU rDNA), two internal transcribed spacer (ITS), and two external transcribed spacer (ETS). From the aforementioned, only ETS fragments have not been used in molecular diagnostic of *Sarcocystis* spp.

Genotyping of SSU rRNA

In 1994, Tenter et al. created a molecular diagnostic test for *S. tenella*, *S. arieticanis*, *S. gigantea*, and *T. gondii* differentiation in sheep. The test was based on ssu rDNA PCR. This type of assay is often used in phylogenetic analysis (Woese et al. 1990, Neefs et al. 1991). This sequence is conservative for related organisms, and has a variable region characteristic for the species (Maidak et al. 1997), which makes it a good diagnostic marker. SSU rDNA sequence was also used in studies based on hybridization techniques with ³²P isotope, in order to diversify *S. cruzi*, *S. tenella*, *S. fusiformis*, *S. gigantea*, and *T. gondii* (Holmdahl et al. 1993). Diagnostics of *Sarcocystis* spp. based on SSU rRNA was performed in cattle (Fischer et al 1998), horses (Elsheikha and Mansfield 2007), and pigs (Caspari et al. 2010). The investigation of Yang et al. (2001) is an example of using SSU rRNA in phylogenetic analysis. Investigators have proved that in both bovine and buffalo (*Bubalus bubalis*), the same species of *Sarcocystis* are present. Oryan et al. (2011) used the SSU rDNA fragment to investigate *S. fusiformis* from water buffalo by means of RFLP and sequencing. Heterogeneity between isolates from various geographical localizations was reported. Other protozoa, from *Apicomplexa* phylum, were studied by the same fragment of rDNA (Morrison and Ellis 1997). This type of investigations were performed in numerous wild animals including pigeons (*Columba livia f. domestica*) (Olias et al. 2009), black bears (*Ursus americanus*) (Davies et al. 2011), raccoon dogs (*Nyctereutes procyonoides*) (Kubo et al. 2010), sparrowhawks (*Accipiter nisus*) (Olias et al. 2010), ducks (*Anas platyrhynchos*) (Kutkienl et al. 2011), roe deers (*Capreolus capreolus*) (Dahlgren and Gjerde 2008), moose (*Alces alces*), red foxes (*Vulpes vulpes*) (Dahlgren and Gjerde 2010), wolverines (*Gulo gulo*) (Dubey et al. 2010), vipers (*Atheris nitschei*) (Slapeta et al. 2003), and otters (*Enhydra lutris nereis*) (Miller et al. 2009).

Genotyping of LSU rRNA

Rbotyping, based on SSU rDNA in some cases, did not provide comprehensive results for scientists.

Usage of large ribosomal subunit (LSU rDNA, 28S rDNA) in *Sarcocystis* spp. diagnostics can enhance the sensitivity of investigation. This sequence is defined as a high variable DNA fragment at the interspecies level. Data obtained from LSU rDNA analysis were useful in sophisticated phylogenetic studies, and verified results of other assays (Mugridge et al. 2000). An example of the survey performed with LSU rDNA is Slapeta et al. (2003) work on the occurrence of *Sarcocystis* in vipers. Investigators used the D2 domain of mentioned nucleic acid fragment, and made phylogenetic parallel. LSU rDNA was used in the diagnosis of *S. rileyi*, found in mallard duck (*Anas platyrhynchos*). It was the first detection of this parasite in Europe (Kutkienl et al. 2011). The same type of sequence has also been used to confirm that *S. wobeseri* is a parasite common in both mallard ducks as well as in Barnacle goose (*Branta leucopsis*) (Kutkienl et al. 2010). LSU rDNA was also used to phenotype *S. cornix* in corvids (Kutkienl et al. 2008). Olias et al. (2010) applied this tool to examine *Sarcocystis* spp. in pigeons (*Columba livia f. domestica*) and sparrowhawks (*Accipiter nisus*), what resulted in detection of two new *Sarcocystis* species.

Genotyping of ITS fragments

Both, a ITS-1 and ITS-2 are useful molecular markers in population genetics studies. These fragments of nucleic acids, are adjacent to conservative genes that encode ribosomal RNA. ITS sequences are determined as species/strain variable (McManus and Bowles 1996). These features had an impact on ITS fragments popularity in genotyping of *T. gondii* and *Neospora* spp. (Homan et al. 1997). It was confirmed by Su et al. (2003). The investigators performed sensitive PCR assay for *Eimeria* spp. derived from poultry. ITS fragments were useful in *Sarcocystis* spp. diagnostic as well. With the use of described genom fragment, Rosenthal et al. (2008) compared *S. cruzi* isolates from North America with isolates from South America. They confirmed that both populations were mixed and *S. cruzi* can cross intercontinental borders.

Interspecies differentiation of *Sarcocystis* taxa, based on ITS fragments is a common practice in researches. An example of this type of application is *S. neurona* and *S. falcatulta* molecular differentiation assay (Marsh et al. 1999). It is also a useful tool in new *Sarcocystis* species/strains searching, because it is a conservative molecular marker (Gozalo et al. 2007). ITS fragments analysis were used in numerous studies concerning wild animals, including pigeons (*Columba livia f. domestica*) (Olias et al. 2009), rhesus macaque (*Macaca mulatta*), ducks (*Anas platyrhynchos*) (Kut-

kienl et al. 2011), otters (*Enhydra lutris nereis*) (Miller et al. 2009), wolverines (*Gulo gulo*) (Dubey et al. 2010), sparrowhawks (*Accipiter nisus*), (Olias et al. 2010), bald eagles (*Haliaeetus leucocephalus*), and golden eagles (*Aquila chrysaetos*) (Wunschmann et al. 2010).

In some cases, additional analysis of nucleic acids is required. For this purpose, scientists can use restriction fragment length polymorphism (RFLP-PCR) and single nucleotide polymorphism (SNP) analyses.

Restriction Fragment Length Polymorphism (RFLP)

An important tool used in the diagnosis of *Sarcocystis* spp. is restriction fragment length polymorphism (RFLP) analysis. This method is based on fragmentation of nucleic acids obtained from PCR reaction. In order to carry out RFLP assay, one must digest nucleic acids by restriction enzymes. After digestion, investigators receive a specific pattern of nucleic acids fragments. Comparison between obtained fragments define sequences variation. The described method is inexpensive, easy to use, and provides high sensitivity of measurements (McManus and Bowles 1996). A RFLP assay was used to examine ITS-1 fragments. In this investigation, *S. cruzi* was determined as infective for cattle and buffalo (Li et al. 2002). Tanhauser et al. (1999) performed RFLP based method that allowed for *S. falcatulta* and *S. neurona* differentiation. Opossum feces were investigated in this study. The described method is also a useful to investigate stool samples from humans, cats and dogs (Xiang et al. 2009). Universality of restriction enzymes creates possibilities in diagnostics of both domestic (More et al. 2011) and wild animals (Kia et al. 2011). Based on this type of assay, molecular profile of *S. cameli* has been described (Motamedi et al. 2011). Yang et al. (2002) summarized the use of RFLP in *Sarcocystis* spp. differentiation in domestic animals. The authors have created a solution of restriction enzymes selection in order to obtain interspecies distinction of parasite.

Single nucleotide polymorphism (SNP)

Nucleotide polymorphism is caused by an evolution – internal agent. Genetic polymorphisms can also be the result of viruses or radiation impact (external agents). SNPs are present in all regions of the non-coding and coding regions of a genomes. Moreover, nucleotide variation may affect amino acid sequence in proteins. SNP analysis is common in mod-

ern evolutionary researches, so that investigators can seek for correlations between different species/strains (Brown 2007). Dahlgren and Gjerde (2008) carried out SNP analysis of six *Sarcocystis* species occurring in moose. *S. hardangeri* was established as genetically most variable among other species.

Whole genome analysis methods and tandem repeats investigations were used in order to receive data about phylogenetic and epidemiological studies of *Sarcocystis* spp.

Random Amplification of Polymorphic DNA (RAPD)

RAPD is a PCR reaction variant. In contrast to classical PCR, in the RAPD method, random primers (10-15 nucleotides) are used. The template for the reaction is whole genomic DNA. This results in amplification of nucleic acid fragments, which are various and specific for the organism examined. The reaction is usually visualized on electrophoresis method by different pattern of bands. RAPD is a proper tool for insertions/deletions detection, gene mapping, gene localization, phylogenetic research, the evolutionary markers detection. This method is time effective, easy-to-use, knowledge about the sequence of DNA is not required, and relatively small amounts of DNA are needed to perform the reaction (Bardakci 2001). The RAPD method was used by MacPherson and Gajadhar (1994) in a molecular probe for *S. cruzi* in a cattle investigation. The aforementioned study inspired other investigators to work out a sensitive *S. cruzi*, *S. hirsuta*, *S. hominis* detection method (Guclu et al. 2004). This type of investigation was used for detecting *Sarcocystis* spp. in sheep (Joachim et al. 1996) and differentiation between *S. neurona* and *S. falcatula* (Tanhauser et al. 1999). In 1994, Granstrom et. al. described a sensitive method for *S. neurona* detection. Investigators obtained 550 bp DNA fragment by means of 16 nucleotides primer. This marker was sensitive for *S. neurona* among 8 coccidia species, inter alia other *Sarcocystis* spp., *T. gondii*, and *Eimeria* spp. This method was also used in the diagnosis of *Sarcocystis* in the domestic cat (Gillis et al. 2003) and brown-headed cowbird (*Molothrus ater*) (Mansfield et al. 2008).

Amplified Fragment Length Polymorphism PCR (AFLP)

Methodology of AFLP PCR comprises several steps. First, restriction enzymes digest the whole genomic DNA, followed by ligation of adaptors to the

sticky ends of the restriction fragments. Reaction is performed by ligase T4 DNA. Subsequently, two PCR reactions should be performed. Primers for PCR are complementary to adapter sequences. Low specific primers are used in the first PCR, and specific are used in the second reaction. Amplification products are visualized on the electrophoretic gel, usually polyacrylamide. This method is characterized by high resolution and repeatability. The knowledge about the sequence of the template is not required. This type of assay gives investigators a genetic fingerprint, which enables diagnostics and genotyping studies (Vos et al. 1995). AFLP was used to analyze *S. neurona* derived from different hosts. *S. neurona* was characterized by high intraspecies genetic variability, from 82% to 94% of whole genomic DNA. Sixty four primers were examined, nine of which gave a high resolution image in relation to polymorphism and phylogeny of *S. neurona* (Elsheikha et al. 2006).

Tandem Repeats Analysis (Microsatellite sequences)

Microsatellite sequences are common in eukaryotic genomes and consist of many repeated 2 to 6 nucleotides fragments. These sequences are highly variable, which is associated with mutation occurrence. The aforementioned feature gives the possibility to make genetic variations and to conduct phylogenetic studies. Usually, the difference between fragments length is investigated. Changes in the nucleotide sequences composition of studied tandem repeats are not very important. Most of the microsatellites occur at the end of chromosomes. These types of genetic markers are used in genotyping, evolutionary studies, and population studies (Ellegren 2004).

The investigation concerning *Sarcocystis* spp. phylogeography, twelve microsatellite markers were analyzed. Samples were collected from distant geographical locations and were investigated in terms of *S. neurona* and *S. falcatula* derived from horses, sea otters and opossums. The occurrence of one genotype was frequently established in all examined species of animals. However, genetic variation was estimated at a high level. *S. neurona* from the North American population proved to be genetically homogeneous (Asmundsson et al. 2006).

Conclusions

The development of molecular biology has contributed to improve diagnostic methods and

phylogenetics. *Sarcocystis* spp. has become one of the targets of molecular parasitologists. Many investigations were inspired by *Sarcocystis* spp. morphological similarity and species richness of the aforementioned family. There are many useful tools for differentiation of *Sarcocystis* spp., which allow scientists to carry out epidemiological studies. However, there is no sequenced genome of any species from *Sarcocystis* family so far. Lack of clarity associated with the pathogenicity of this protozoan, the growing number of diagnosed sarcocystosis cases, and the zoonotic potential of *Sarcocystis* spp. encourage researchers to make further studies on this group of organisms.

References

- Abdul-Rahman SM, Rashad SM, Doma MA (2002) Human muscle sarcocystosis in relation to non-specific rheumatic diseases and rheumatoid arthritis. *Egyptian Rheumatology and Rehabilitation* 29: 743-753.
- Acha PN, Szyfres B (2003) Zoonoses and communicable diseases common to man and animals, 3rd ed., vol. III: Parasitoses, Pan American Health Organization (PAHO), Washington D.C.
- Arness MK, Brown JD, Dubey JP, Neafie RC, Granstrom DE (1999) An outbreak of acute eosinophilic myositis attributed to human *Sarcocystis* parasitism. *Am J Trop Med Hyg* 61: 548-553.
- Asmundsson IM, Dubey JP, Rosenthal BM (2006) A genetically diverse but distinct North American population of *Sarcocystis neurona* includes an overrepresented clone described by 12 microsatellite alleles. *Infect Genet Evol* 6: 352-360.
- Bardakci F (2001) Random amplified polymorphic DNA (RAPD) markers. *Turkish Journal of Biology* 25: 185-196.
- Brown TA (2007) *Genomes*, 3rd ed., Garland Science, New York.
- Caspari K, Grimm F, Kühn N, Caspari NC, Basso W (2010) First report of naturally acquired clinical sarcocystosis in a pig breeding stock. *Vet Parasitol* 177: 175-178.
- Dahlgren SS, Gjerde B (2008) *Sarcocystis* in moose (*Alces alces*): molecular identification and phylogeny of six *Sarcocystis* species in moose, and a morphological description of three new species. *Parasitol Res* 103: 93-110.
- Dahlgren SS, Gjerde B (2010) The red fox (*Vulpes vulpes*) and the arctic fox (*Vulpes lagopus*) are definitive hosts of *Sarcocystis alces* and *Sarcocystis hjorti* from moose (*Alces alces*). *Parasitology* 137: 1547-1557.
- Davies JL, Haldorson GJ, Bradway DS, Britton AP (2011) Fatal hepatic sarcocystosis in a captive black bear (*Ursus americanus*) associated with *Sarcocystis canis*-like infection. *J Vet Diagn Invest* 23: 379-383.
- Dubey JP, Reichard MV, Torretti L, Garvon JM, Sundar N, Grigg ME (2010) Two new species of *Sarcocystis* (Apicomplexa: Sarcocystidae) infecting the wolverine (*Gulo gulo*) from Nunavut, Canada. *J Parasitol* 96: 972-976.
- Dubey JP, Speer CA, Charleston WA (1989a) Ultrastructural differentiation between sarcocysts of *Sarcocystis hirsuta* and *Sarcocystis hominis*. *Vet Parasitol* 34: 153-157.
- Dubey JP, Speer CA, Fayer R (1989b) *Sarcocystosis of animals and man*, 1st ed., CRC Press, Boca Raton.
- Ellegren H (2004) Microsatellites: simple sequences with complex evolution. *Nat Rev Genet* 5: 435-445.
- Elsheikha HM, Mansfield LS (2007) Molecular typing of *Sarcocystis neurona*: current status and future trends. *Vet Parasitol* 149: 43-55.
- Elsheikha HM, Schott HC 2nd, Mansfield LS (2006) Genetic variation among isolates of *Sarcocystis neurona*, the agent of protozoal myeloencephalitis, as revealed by amplified fragment length polymorphism markers. *Infect Immun* 74: 3448-3454.
- Fayer R (2004) *Sarcocystis* spp. in Human Infections. *Clin Microbiol Rev* 17: 894-902.
- Fischer S, Odening K (1998) Characterization of bovine *Sarcocystis* species by analysis of their 18S ribosomal DNA sequences. *J Parasitol* 84: 50-54.
- Giboda M, Ditrich O, Schoz T, Viengsay T, Bouaphanh S (1991) Current status of food-borne parasitic zoonoses in Laos. *Southeast Asian J Trop Med Public Health* 22: 56-61.
- Gillis KD, MacKay RJ, Yowell CA, Levy JK, Greiner EC, Dame JB, Cheadle MA, Hernandez J, Massey ET (2003) Naturally occurring *Sarcocystis* infection in domestic cats (*Felis catus*). *Int J Parasitol* 33: 877-883.
- Gozal AS, Montali RJ, St Claire M, Barr B, Rejmanek D, Ward JM (2007) Chronic polymyositis associated with disseminated sarcocystosis in a captive-born rhesus macaque. *Vet Pathol* 44: 695-699.
- Granstrom DE, MacPherson JM, Gajadhar AA, Dubey JP, Tramontin R, Stamper S (1994) Differentiation of *Sarcocystis neurona* from eight related coccidia by random amplified polymorphic DNA assay. *Mol Cell Probes* 8: 353-356.
- Greve E (1985) Sarcocystosis--an overlooked zoonosis. Man as intermediate and final host. *Dan Med Bull* 32: 228-230.
- Guclu F, Aldem OS, Guler L (2004) Differential identification of cattle *Sarcocystis* spp. by random amplified Polymorphic DNA Polymerase chain reaction (RAPD-PCR). *Revue d Elevage et de Medecine Veterynaire* 155: 440-444.
- Habeeb YS, Selim MA, Ali MS, Mahmoud LA, Abdel Hadi AM, Shafei A (1996) Serological diagnosis of extraintestinal Sarcocystosis. *J Egypt Soc Parasitol* 26: 393-400.
- Heckerroth AR, Tenter AM (1999) Comparison of immunological and molecular methods for the diagnosis of infections with pathogenic *Sarcocystis* species in sheep. *Toikai J Exp Clin Med* 23: 293-302.
- Holmdahl OJ, Mattsson JG, Uggla A, Johansson KE (1993) Oligonucleotide probes complementary to variable regions of 18S rRNA from *Sarcocystis* species. *Mol Cell Probes* 7: 481-486.
- Homan WL, Limper L, Verlaan M, Borst A, Vercammen M, van Knapen F (1997) Comparison of the internal transcribed spacer, ITS 1, from *Toxoplasma gondii* isolates and *Neospora caninum*. *Parasitol Res* 83: 285-289.
- Jehle C, Dinkel A, Sander A, Morent M, Romig T, Luc PV, De TV, Thai VV, Mackenstedt U (2009) Diagnosis of *Sarcocystis* spp. in cattle (*Bos taurus*) and water buffalo (*Bubalus bubalis*) in Northern Vietnam. *Vet Parasitol* 166: 314-320.
- Joachim A, Tenter AM, Jeffries AC, Johnson AM (1996) A RAPD-PCR derived marker can differentiate between

- pathogenic and non-pathogenic *Sarcocystis* species of sheep. *Mol Cell Probes* 10: 165-172.
- Kia EB, Mirhendi H, Rezaeian M, Zahabiun F, Sharbatkhori M (2011) First molecular identification of *Sarcocystis miescheriana* (Protozoa, Apicomplexa) from wild boar (*Sus scrofa*) in Iran. *Exp Parasitol* 127: 724-726.
- Kubo M, Kawachi T, Murakami M, Kubo M, Tokuhiko S, Agatsuma T, Ito K, Okano T, Asano M, Fukushi H, Nagataki M, Sakai H, Yanai T (2010) Meningoencephalitis associated with *Sarcocystis* spp. in a free-living Japanese raccoon dog (*Nyctereutes procyonoides viverrinus*). *J Comp Pathol* 143: 185-189.
- Kutkienė L, Prakas P, Sruoga A, Butkauskas D (2008) *Sarcocystis* in the birds family Corvidae with description of *Sarcocystis cornix* sp. nov. from the hooded crow (*Corvus cornix*). *Parasitol Res* 104: 329-336.
- Kutkienė L, Prakas P, Sruoga A, Butkauskas D (2010) The mallard duck (*Anas platyrhynchos*) as intermediate host for *Sarcocystis wobeseri* sp. nov. from the barnacle goose (*Branta leucopsis*). *Parasitol Res* 107: 879-888.
- Kutkienė L, Prakas P, Sruoga A, Butkauskas D (2011) Identification of *Sarcocystis rileyi* from the mallard duck (*Anas platyrhynchos*) in Europe: cyst morphology and results of DNA analysis. *Parasitol Res* 108: 709-714.
- Li QQ, Yang ZQ, Zuo YX, Attwood SW, Chen XW, Zhang YP (2002) A PCR-based RFLP analysis of *Sarcocystis cruzi* (Protozoa: Sarcocystidae) in Yunnan Province, PR China, reveals the water buffalo (*Bubalus bubalis*) as a natural intermediate host. *J Parasitol* 88: 1259-1261.
- MacPherson JM, Gajadhar AA (1994) Specific amplification of *Sarcocystis cruzi* DNA using a randomly primed polymerase chain reaction assay. *Vet Parasitol* 55: 267-277.
- Maidak BL, Olsen GJ, Larsen N, Overbeek R, McCaughey MJ, Woese CR (1997) The RDP (Ribosomal Database Project). *Nucleic Acids Res* 25: 109-111.
- Mansfield LS, Mehler S, Nelson K, Elsheikha HM, Murphy AJ, Knust B, Tanhauser SM, Gearhart PM, Rossano MG, Bowman DD, Schott HC, Patterson JS (2008) Brown-headed cowbirds (*Molothrus ater*) harbour *Sarcocystis neurona* and act as intermediate hosts. *Vet Parasitol* 153: 24-43.
- Marsh AE, Barr BC, Tell L, Bowman DD, Conrad PA, Ketcherside C, Green T (1999) Comparison of the internal transcribed spacer, ITS-1, from *Sarcocystis falcatula* isolates and *Sarcocystis neurona*. *J Parasitol* 85: 750-757.
- McManus DP, Bowles J (1996) Molecular genetic approaches to parasite identification: their value in diagnostic parasitology and systematics. *Int J Parasitol* 26: 687-704.
- Miller MA, Barr BC, Nordhausen R, James ER, Magargal SJ, Murray M, Conrad PA, Toy-Choutka S, Jessup DA, Grigg ME (2009) Ultrastructural and molecular confirmation of the development of *Sarcocystis neurona* tissue cysts in the central nervous system of southern sea otters (*Enhydra lutris nereis*). *Int J Parasitol* 39: 1363-1372.
- Moré G, Abrahamovich P, Jurado S, Bacigalupe D, Marin JC, Rambeaud M, Venturini L, Venturini MC (2011) Prevalence of *Sarcocystis* spp. in Argentinean cattle. *Vet Parasitol* 177: 162-165.
- Morrison DA, Ellis JT (1997) Effects of nucleotide sequence alignment on phylogeny estimation: a case study of 18S rDNAs of apicomplexa. *Mol Biol Evol* 14: 428-441.
- Motamedi GR, Dalimi A, Nouri A, Aghaeipour K (2011) Ultrastructural and molecular characterization of *Sarcocystis* isolated from camel (*Camelus dromedarius*) in Iran. *Parasitol Res* 108: 949-954.
- Mugridge NB, Morrison DA, Jäkel T, Heckerroth AR, Tenter AM, Johnson AM (2000) Effects of sequence alignment and structural domains of ribosomal DNA on phylogeny reconstruction for the protozoan family Sarcocystidae. *Mol Biol Evol* 17: 1842-1853.
- Neefs JM, Van de Peer Y, De Rijk P, Goris A, De Wachter R (1991) Compilation of small ribosomal subunit RNA sequences. *Nucleic Acids Res* 19: 1987-2015.
- Olias P, Gruber AD, Heydorn AO, Kohls A, Mehlhorn H, Hafez HM, Lierz M (2009) A novel *Sarcocystis*-associated encephalitis and myositis in racing pigeons. *Avian Pathol* 38: 121-128.
- Olias P, Olias L, Lierz M, Mehlhorn H, Gruber AD (2010) *Sarcocystis calchasi* is distinct to *Sarcocystis columbae* sp. nov. from the wood pigeon (*Columba palumbus*) and *Sarcocystis* sp. from the sparrowhawk (*Accipiter nisus*). *Vet Parasitol* 171: 7-14.
- Oryan A, Sharifiyazdi H, Khordadmehr M, Larki S (2011) Characterization of *Sarcocystis fusiformis* based on sequencing and PCR-RFLP in water buffalo (*Bubalus bubalis*) in Iran. *Parasitol Res* 109: 1563-1570.
- Pamphlett R, O'Donoghue P (1990) *Sarcocystis* infection of human muscle. *Aust N Z J Med* 20: 705-707.
- Pritt B, Trainer T, Simmons-Arnold L, Evans M, Dunams D, Rosenthal BM (2008) Detection of *Sarcocystis* parasites in retail beef: a regional survey combining histological and genetic detection methods. *J Food Prot* 71: 2144-2147.
- Reiner G, Hübner K, Hepp S (2009) Suffering in diseased pigs as expressed by behavioural, clinical and clinical-chemical traits, in a well defined parasite model. *Appl Anim Behav Sci* 118: 222-231.
- Rosenthal BM, Dunams DB, Pritt B (2008) Restricted genetic diversity in the ubiquitous cattle parasite, *Sarcocystis cruzi*. *Infect Genet Evol* 8: 588-592.
- Saleque A, Bhatia BB (1991) Prevalence of *Sarcocystis* in domestic pigs in India. *Vet Parasitol* 40: 151-153.
- Slapeta JR, Modrý D, Votýpka J, Jirkú M, Lukes J, Koudela B (2003) Evolutionary relationships among cyst-forming coccidia *Sarcocystis* spp. (Alveolata: Apicomplexa: Coccidia) in endemic African tree vipers and perspective for evolution of heteroxenous life cycle. *Mol Phylogenet Evol* 27: 464-475.
- Su YC, Fei AC, Tsai FM (2003) Differential diagnosis of five avian *Eimeria* species by polymerase chain reaction using primers derived from the internal transcribed spacer 1 (ITS-1) sequence. *Vet Parasitol* 117: 221-227.
- Tanhauser SM, Yowell CA, Cutler TJ, Greiner EC, MacKay RJ, Dame JB (1999) Multiple DNA markers differentiate *Sarcocystis neurona* and *Sarcocystis falcatula*. *J Parasitol* 85: 221-228.
- Tenter AM (1995) Current research on *Sarcocystis* species of domestic animals. *Int J Parasitol* 25: 1311-1330.
- Tenter AM, Luton K, Johnson AM (1994) Species specific identification of *Sarcocystis* and *Toxoplasma* by PCR amplification of small subunit ribosomal RNA gene fragments. *Appl Parasitol* 35: 173-188.
- Tenter AM, Vietmeyer C, Thummel P, Rommel M (1991) Detection of species-specific and cross-reactive epitopes in *Sarcocystis* cystozoites by monoclonal antibodies. *Parasitol Res* 77: 212-216.
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M,

- Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23: 4407-4414.
- Wilairatana P, Radomyos P, Radomyos B, Phraevanich R, Plooksawasdi W, Chanthavanich P, Viravan C, Looareesuwan S (1996) Intestinal sarcocystosis in Thai laborers. *Southeast Asian J Trop Med Public Health* 27: 43-46.
- Woese CR, Kandler O, Wheelis ML (1990) Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci USA* 87: 4576-4579.
- Wong KT, Pathmanathan R (1992) High prevalence of human skeletal muscle sarcocystosis in south-east Asia. *Trans R Soc Trop Med Hyg* 86: 631-632.
- World Health Organization (1981) Intestinal protozoan and helminthic infections, vol. 666. World Health Organization, Geneva.
- Wunschmann A, Rejmanek D, Conrad PA, Hall N, Cruz-Martinez L, Vaughn SB, Barr BC (2010) Natural fatal *Sarcocystis falcatula* infections in free-ranging eagles in North America. *J Vet Diagn Invest* 22: 282-289.
- Xiang Z, Chen X, Yang L, He Y, Jiang R, Rosenthal BM, Luan P, Attwood SW, Zuo Y, Zhang YP, Yang Z (2009) Non-invasive methods for identifying oocysts of *Sarcocystis* spp. from definitive hosts. *Parasitol Int* 58: 293-296.
- Yang ZQ, Li QQ, Zuo YX, Chen XW, Chen YJ, Nie L, Wei CG, Zen JS, Attwood SW, Zhang XZ, Zhang YP (2002) Characterization of *Sarcocystis* species in domestic animals using a PCR-RLFP analysis of variation in the 18S rRNA gene: a cost-effective and simple technique for routine species identification. *Exp Parasitol* 102: 212-217.
- Yang ZQ, Zuo YX, Yao YG, Chen XW, Yang GC, Zhang YP (2001). Analysis of the 18S rRNA genes of *Sarcocystis* species suggests that the morphologically similar organisms from cattle and water buffalo should be considered the same species. *Mol Biochem Parasitol* 115: 283-288.